

Talk to your child's doctor or nurse about the vaccines recommended for their age.

	Flu <i>Influenza</i>	Tdap Tetanus, diphtheria, pertussis	HPV Human papillomavirus	Meningococcal		Pneumococcal	Hepatitis B	Hepatitis A		Inactivated Polio	MMR Measles, mumps, rubella	Chickenpox <i>Varicella</i>
				MenACWY	MenB							
7-8 Years	Green	Orange		Orange		Orange	Orange	Orange	Orange	Orange	Orange	Orange
9-10 Years	Green	Orange	Green, Blue	Orange		Orange	Orange	Orange	Orange	Orange	Orange	Orange
11-12 Years	Green	Green, Orange	Green	Green, Orange		Orange	Orange	Orange	Orange	Orange	Orange	Orange
13-15 Years	Green	Orange	Orange	Orange		Orange	Orange	Orange	Orange	Orange	Orange	Orange
16-18 Years	Green	Orange	Orange	Green, Orange		Orange	Orange	Orange	Orange	Orange	Orange	Orange

More information:


Preteens and teens should get a flu vaccine every year.


Preteens and teens should get one shot of Tdap at age 11 or 12 years.


All 11-12 year olds should get a 2-shot series of HPV vaccine at least 6 months apart. A 3-shot series is needed for those with weakened immune systems and those age 15 or older.


All 11-12 year olds should get a single shot of a quadrivalent meningococcal conjugate vaccine (MenACWY). A booster shot is recommended at age 16.

Teens, 16-18 years old, **may** be vaccinated with a MenB vaccine.

 These shaded boxes indicate when the vaccine is recommended for all children unless your doctor tells you that your child cannot safely receive the vaccine.

 These shaded boxes indicate the vaccine should be given if a child is catching-up on missed vaccines.

 These shaded boxes indicate the vaccine is recommended for children with certain health or lifestyle conditions that put them at an increased risk for serious diseases. See vaccine-specific recommendations at www.cdc.gov/vaccines/pubs/ACIP-list.htm.

 This shaded box indicates the vaccine is recommended for children not at increased risk but who wish to get the vaccine after speaking to a provider.



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention

American Academy of Pediatrics
DEDICATED TO THE HEALTH OF ALL CHILDREN™



AMERICAN ACADEMY OF FAMILY PHYSICIANS
STRONG MEDICINE FOR AMERICA

Vaccine-Preventable Diseases and the Vaccines that Prevent Them

Diphtheria (Can be prevented by Tdap vaccination)

Diphtheria is a very contagious bacterial disease that affects the respiratory system, including the lungs. Diphtheria bacteria can be passed from person to person by direct contact with droplets from an infected person's cough or sneeze. When people are infected, the bacteria can produce a toxin (poison) in the body that can cause a thick coating in the back of the nose or throat that makes it hard to breathe or swallow. Effects from this toxin can also lead to swelling of the heart muscle and, in some cases, heart failure. In serious cases, the illness can cause coma, paralysis, and even death.

Hepatitis A (Can be prevented by HepA vaccination)

Hepatitis A is an infection in the liver caused by hepatitis A virus. The virus is spread primarily person-to-person through the fecal-oral route. In other words, the virus is taken in by mouth from contact with objects, food, or drinks contaminated by the feces (stool) of an infected person. Symptoms can include fever, tiredness, poor appetite, vomiting, stomach pain, and sometimes jaundice (when skin and eyes turn yellow). An infected person may have no symptoms, may have mild illness for a week or two, may have severe illness for several months, or may rarely develop liver failure and die from the infection. In the U.S., about 100 people a year die from hepatitis A.

Hepatitis B (Can be prevented by HepB vaccination)

Hepatitis B causes a flu-like illness with loss of appetite, nausea, vomiting, rashes, joint pain, and jaundice. Symptoms of acute hepatitis B include fever, fatigue, loss of appetite, nausea, vomiting, pain in joints and stomach, dark urine, grey-colored stools, and jaundice (when skin and eyes turn yellow).

Human Papillomavirus (Can be prevented by HPV vaccination)

Human papillomavirus is a common virus. HPV is most common in people in their teens and early 20s. It is the major cause of cervical cancer in women and genital warts in women and men. The strains of HPV that cause cervical cancer and genital warts are spread during sex.

Influenza (Can be prevented by annual flu vaccination)

Influenza is a highly contagious viral infection of the nose, throat, and lungs. The virus spreads easily through droplets when an infected person coughs or sneezes and can cause mild to severe illness. Typical symptoms include a sudden high fever, chills, a dry cough, headache, runny nose, sore throat, and muscle and joint pain. Extreme fatigue can last from several days to weeks. Influenza may lead to hospitalization or even death, even among previously healthy children.

Measles (Can be prevented by MMR vaccination)

Measles is one of the most contagious viral diseases. Measles virus is spread by direct contact with the airborne respiratory droplets of an infected person. Measles is so contagious that just

being in the same room after a person who has measles has already left can result in infection. Symptoms usually include a rash, fever, cough, and red, watery eyes. Fever can persist, rash can last for up to a week, and coughing can last about 10 days. Measles can also cause pneumonia, seizures, brain damage, or death.

Meningococcal Disease (Can be prevented by meningococcal vaccination)

Meningococcal disease is caused by bacteria and is a leading cause of bacterial meningitis (infection around the brain and spinal cord) in children. The bacteria are spread through the exchange of nose and throat droplets, such as when coughing, sneezing or kissing. Symptoms include sudden onset of fever, headache, and stiff neck. Meningococcal bacteria also cause blood infections. About one of every ten people who get the disease dies from it. Survivors of meningococcal disease may lose their arms or legs, become deaf, have problems with their nervous systems, become developmentally disabled, or suffer seizures or strokes.

Mumps (Can be prevented by MMR vaccination)

Mumps is an infectious disease caused by the mumps virus, which is spread in the air by a cough or sneeze from an infected person. A child can also get infected with mumps by coming in contact with a contaminated object, like a toy. The mumps virus causes swollen salivary glands under the ears or jaw, fever, muscle aches, tiredness, abdominal pain, and loss of appetite. Severe complications for children who get mumps are uncommon, but can include meningitis (infection of the covering of the brain and spinal cord), encephalitis (inflammation of the brain), permanent hearing loss, or swelling of the testes, which rarely results in decreased fertility.

Pertussis (Whooping Cough) (Can be prevented by Tdap vaccination)

Pertussis is caused by bacteria spread through direct contact with respiratory droplets when an infected person coughs or sneezes. In the beginning, symptoms of pertussis are similar to the common cold, including runny nose, sneezing, and cough. After 1-2 weeks, pertussis can cause spells of violent coughing and choking, making it hard to breathe, drink, or eat. This cough can last for weeks. Pertussis is most serious for babies, who can get pneumonia, have seizures, become brain damaged, or even die. About half of children under 1 year of age who get pertussis must be hospitalized.

Pneumococcal Disease (Can be prevented by pneumococcal vaccination)

Pneumonia is an infection of the lungs that can be caused by the bacteria called pneumococcus. This bacteria can cause other types of infections too, such as ear infections, sinus infections, meningitis (infection of the covering around the brain and spinal cord), and bacteremia (bloodstream infection). Sinus and ear infections are usually mild and are much more common than the

more serious forms of pneumococcal disease. However, in some cases pneumococcal disease can be fatal or result in long-term problems, like brain damage and hearing loss. Pneumococcal disease spreads when people cough or sneeze. Many people have the bacteria in their nose or throat at one time or another without being ill—this is known as being a carrier.

Polio (Can be prevented by IPV vaccination)

Polio is caused by a virus that lives in an infected person's throat and intestines. It spreads through contact with the stool of an infected person and through droplets from a sneeze or cough. Symptoms typically include sore throat, fever, tiredness, nausea, headache, or stomach pain. In about 1% of cases, polio can cause paralysis. Among those who are paralyzed, About 2 to 10 children out of 100 die because the virus affects the muscles that help them breathe.

Rubella (German Measles) (Can be prevented by MMR vaccination)

Rubella is caused by a virus that is spread through coughing and sneezing. In children rubella usually causes a mild illness with fever, swollen glands, and a rash that lasts about 3 days. Rubella rarely causes serious illness or complications in children, but can be very serious to a baby in the womb. If a pregnant woman is infected, the result to the baby can be devastating, including miscarriage, serious heart defects, mental retardation and loss of hearing and eye sight.

Tetanus (Lockjaw) (Can be prevented by Tdap vaccination)

Tetanus is caused by bacteria found in soil, dust, and manure. The bacteria enters the body through a puncture, cut, or sore on the skin. When people are infected, the bacteria produce a toxin (poison) that causes muscles to become tight, which is very painful. Tetanus mainly affects the neck and belly. This can lead to "locking" of the jaw so a person cannot open his or her mouth, swallow, or breathe. Complete recovery from tetanus can take months. One to two out of 10 people who get tetanus die from the disease.

Varicella (Chickenpox) (Can be prevented by varicella vaccination)

Chickenpox is caused by the varicella zoster virus. Chickenpox is very contagious and spreads very easily from infected people. The virus can spread from either a cough, sneeze. It can also spread from the blisters on the skin, either by touching them or by breathing in these viral particles. Typical symptoms of chickenpox include an itchy rash with blisters, tiredness, headache and fever. Chickenpox is usually mild, but it can lead to severe skin infections, pneumonia, encephalitis (brain swelling), or even death.

If you have any questions about your child's vaccines, talk to your healthcare provider.