



**Summit County Public Health
Influenza Surveillance Report
2022 – 2023 Season
Report #3**



**Flu Surveillance Weeks 3 & 4 (10/16/2022 to 10/29/2022)
Centers for Disease Control and Prevention MMWR Weeks 42 & 43**

Summit County Surveillance Data:

In **Weeks 3 & 4** of influenza surveillance, influenza-related activity was minimal in Summit County.

Table 1: Overall Influenza Activity Indicators in Summit County by week				
	Week 3 MMWR 42 N (%) ¹	Week 4 MMWR 43 N (%) ¹	Percent change from previous week	Number of weeks increasing or decreasing
Lab Reports: Influenza				
Test Performed	964	973	0.9%	↑3
Positive Tests (Number and %)	33 (3.4)	46	38.1%	↑3
Influenza A (Number and %)	31 (3.2)	46	47.0%	↑3
Influenza B (Number and %)	2 (0.2)	0	-100%	↓1
Acute care hospitalizations for Influenza:	2	2	0%	-
Schools absenteeism²	9.0	9.3	2.9%	↑2
Deaths (occurred in Summit County)				
Pneumonia associated	3	1	-66.7%	↓3
Influenza associated	0	0	--	--
COVID-19 associated	7	5	-28.6%	↓2
Emergency room visits (EpiCenter)³ (Figure 3)**				
Total ED Visits	4659	4719	1.3%	↑1
Constitutional Complaints	602 (12.9)	634 (13.4)	4.0%	↑1
Fever and ILI	163 (3.5)	189 (4.0)	14.5%	↑1
1) N and % are reported when available				
2) Absence is for any reason. Percent is from total number of students enrolled. Data was collected from approx. 9 schools or school districts throughout Summit County (n = approx. 32,000 students)				
3)** Percent is from total number of emergency room interactions – elimination of data from a significant reporting facility has resulted in decreases in current and previous week data. Notable changes in EpiCenter data are the result of a temporary programming issue in one of the reporting facilities. Week 3 data has been retroactively adjusted for accurate week-to-week comparison. However, a significant number of ER visits are expected to be unaccounted for at this time.				
Note: Data is provisional and may be updated as more information is received. Percentages should be interpreted with caution. Small changes in number can result in large changes in percent. When a percentage, or prevalence, is available in this table, the percent change will be calculated from those values				

Lab reports: During week 3 and 4 of influenza surveillance, reporting Summit County facilities performed 1,937 flu tests, of which 79 had positive results. **(Figure 4) Note: Influenza data are collected from selected reporting partners and do not represent positivity rates for the entire county.**


Acute Care Hospitalizations: There were two reported admissions during week 3 and week 4. **Figure 2** displays hospitalizations in Summit County.

School absenteeism includes absences regardless of reasoning. In week 3, the absence rate was 9.0% and in week 4 the rate increased to 9.3%.

Zero deaths related to influenza, 5 COVID-19 deaths and 1 pneumonia related deaths occurred in Summit County during Weeks 3 & 4. The number of pneumonia associated deaths and COVID-19 associated deaths decreased in Week 4.


Figure 1 displays weekly counts of deaths occurring in Summit County associated with pneumonia, COVID-19 and influenza.

Figure 1. Weekly Summit County death counts associated with pneumonia and influenza during 2022-2023 season




Hospitalizations: In Week 2, Summit County hospitals reported one influenza-associated hospitalization. In Week 3 there were 2 influenza-associated hospitalizations. **Figure 2** displays weekly confirmed hospitalization counts for Summit County.

Figure 2. Summit County weekly influenza-associated hospitalizations, 2022-2023 season and previous five seasons




EpiCenter collects and analyzes health related data in real time to provide information about the health of the community. This system tracks ER visits related to constitutional complaints and fever and ILI. **Figures 3** displays the weekly number of ER visits related to ILI and flu symptoms in Summit County. There were 184 ILI-related visits reported during Week 3, which was 2.6% of total ED visits (n = 7105). This rate was 5.2% higher than the ILI rate during Week 2.

Figure 3. Weekly ED visits in Summit County related to Fever + ILI stratified by age groups, 2022 to 2023 season**



**Notable changes in EpiCenter data are the result of a temporary programming issue in one of the reporting facilities. Data has been retroactively adjusted for accurate week-to-week comparison. However, a significant number of ER visits are expected to be unaccounted for at this time..

Figure 4. Influenza diagnostic tests with positive results completed by Summit County health facilities, 2022 - 2023 season



Ohio Influenza Activity: from the Ohio Department of Health:

Current Ohio Activity Level (Geographic Spread) – Moderate

During MMWR Week 43, public health surveillance data sources indicate moderate intensity for influenza-like illness (ILI) in outpatient settings reported by Ohio’s sentinel ILINet providers. The percentage of emergency department visits with patients exhibiting constitutional symptoms and Fever/ILI specified ED visits increased and are above baseline levels statewide. Reported cases of influenza-associated hospitalizations increased. There were 135 influenza associated hospitalizations reported during MMWR Week 43.

Data Source	Current week value	Percent Change from last week ¹	# of weeks ²	Trend Chart ³
Influenza-like Illness (ILI) Outpatient Data (ILINet Sentinel Provider Visits)	4.58%	23.78%	↑5	
Thermometer Sales (National Retail Data Monitor) ⁴	0.54%	8.00%	↑2	
Fever and ILI Specified ED Visits (EpiCenter)	2.86%	19.17%	↑5	
Constitutional ED Visits (EpiCenter)	11.79%	10.29%	↑5	
Confirmed Influenza-associated Hospitalizations (Ohio Disease Reporting System)	135	335.48%	↑5	

¹Interpret percent changes with caution. Large variability may be exhibited in data sources with low weekly values.

²Number of weeks that the % change is increasing or decreasing.

³Black lines represent current week’s data; red lines represent baseline averages. The 2020-2021 influenza season has been omitted from the five-year baseline averages due to abnormal counts reported during the COVID-19 pandemic. A five-year average, which includes data from the 2016-2017 season through the 2021-2022 season, is displayed.

⁴Due to abnormally high thermometer sales during the COVID-19 pandemic, the 2019-2020 and 2020-2021 season data has been omitted. A 5-year average, which includes data from the 2015-2016 season through the 2021-2022 season, is shown.

Ohio Influenza Activity Summary Dashboard (October 23 – 29, 2022):

Source <https://odh.ohio.gov/know-our-programs/seasonal-influenza/activity-reports-2022-2023/seasonal-influenza-week-43-20222023>

Ohio Surveillance Data:

- The U.S. World Health Organization (WHO) Collaborating Laboratories System and the National Respiratory and Enteric Virus Surveillance System (NREVSS) has reported 11,401 tests for influenza performed at participating facilities; of these, 32 tested positive for influenza A(H1N1pdm09), 108 for influenza A(H3N2), 206 for influenza A (subtyping not performed), and 15 for influenza B (through 10/30/2022).
- No pediatric influenza-associated mortalities have been reported so far during the 2022-2023 influenza season (through 10/30/2022).
- No novel influenza A virus infections have been reported so far during the 2022-2023 influenza season (through 10/30/2022).
- Incidence of confirmed influenza-associated hospitalizations in 2022-2023 season = 211 (through 10/30/2022).

National Surveillance: from Centers for Disease Control and Prevention (CDC):

National Outpatient Illness Surveillance:

Nationwide during week 43, 4.3% of patient visits reported through ILINet were due to respiratory illness that included fever plus a cough or sore throat, also referred to as ILI. This is an increase compared to week 42 and is above the national baseline of 2.5%. Regions 1, 2, 3, 4, 5, 6, 7, and 9 are above their region-specific baselines. The remaining two regions are below their region-specific baselines. Multiple respiratory viruses are co-circulating, and the relative contribution of influenza virus infection to ILI varies by location.

Figure 5. Percentage of Outpatient Visits for Respiratory Illness reported By the U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet), Weekly National Summary, 2022-2023* and Selected Previous Seasons.





Figure 6. Influenza-like illness (ILI) activity level indicator determined by data reported to ILINet



Source: <https://www.cdc.gov/flu/weekly/>

Global Surveillance:


Influenza Update N° 431, World Health Organization (WHO), published 10/31/2022, based on data up to 10/16/2021. The Update is published every two weeks.

Summary

- Countries are recommended to monitor the co-circulation of influenza and SARS-CoV-2 viruses. They are encouraged to enhance integrated surveillance, and in northern hemisphere countries step-up their influenza vaccination campaign to prevent severe disease and hospitalizations associated with influenza. Clinicians should consider influenza in differential diagnosis, especially for high-risk groups for influenza, and test and treat according to national guidance.
- Globally, influenza activity remained low and where subtyped, influenza A(H3N2) viruses predominated. An increasing trend of influenza activity was observed in the northern hemisphere while a plateau was observed in the southern hemisphere.
- In the countries of North America, influenza activity increased slightly in recent weeks. Influenza A(H3N2) was predominant among the few subtyped viruses.
- In Europe, overall influenza activity remained at inter-seasonal levels, with a low but increasing trend. Influenza A viruses predominated among the reported detections in general with A(H3N2) viruses accounting for the majority of subtyped influenza A viruses.
- In central Asia, Kazakhstan reported increased influenza activity with B/Victoria-lineage viruses predominating.
- In East Asia, influenza activity of predominantly influenza A(H3N2) remained stable, at low levels, overall.
- In Western Asia influenza activity was elevated. Detections of influenza continued to increase in some countries of the Arab Peninsula.

- In the Caribbean and Central American countries, low influenza activity was reported with influenza A(H3N2) most frequently detected.
- In the tropical countries of South America, influenza detections were low and A(H3N2) detections predominated.
- In tropical Africa, influenza activity remained low with detections of influenza A(H3N2), B/Victoria and A(H1N1)pdm09 reported .
- In Southern Asia, influenza activity remained low, with the exception of Iran (Islamic Republic of) where increased activity was reported. The majority of subtyped detections were influenza A(H3N2) and A(H1N1)pdm09, with few influenza B detections.
- In the temperate zones of the southern hemisphere, overall influenza activity appeared to further decrease this reporting period, except in temperate South America where activity increased in some countries.
- In Oceania, influenza activity remained low with detections of influenza A(H1N1)pdm09 and influenza A(H3N2) and some B viruses in Australia. ILI activity in New Zealand and, in general, across the Pacific Islands remained low except in a few countries.

Figure 7. Percentage of respiratory specimens that tested positive for influenza, by influenza transmission zone. Map generated on 28 October 2022.



The boundaries and names shown and the designations used on this map do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted and dashed lines on maps represent approximate border lines for which there may not yet be full agreement.



Data source: Global Influenza Surveillance and Response System (GISRS), FluNet (www.who.int/flu-net)
 Copyright WHO 2022. All rights reserved.

Source: https://cdn.who.int/media/docs/default-source/influenza/influenza-updates/2022/2022_10_31_surveillance_update_431.pdf?sfvrsn=990ecbaa_3&download=true

About this report: Reporting agencies include labs, hospitals, long-term care and community-based care providers, physician offices, university clinic, pharmacies, and schools. Agencies are distributed throughout Summit County and report different indicators of flu activity including total lab tests, numbers of positive tests and type, antiviral prescriptions filled, school absences, and influenza like illness (ILI). Hospitalizations are lab confirmed for influenza and are obtained from the Ohio Disease Reporting System. Number of deaths associated with influenza and pneumonia are gathered from the Summit County Office of Vital Records death listings. Emergency room visits for complaints related to influenza are obtained by syndromic surveillance system (Epicenter).

Special thanks to all agencies who report Influenza related data weekly.

Reporting from participants may not be complete each week. Numbers may change as updated reports are received. For questions, please contact Julie Zidones at the Summit County Public Health Communicable Disease Unit (330-375-2662 or cdu@schd.org). This report was issued on November 4, 2022.